

# WELCOME

# TO



# FDSPharmacy

Learn and Educate

This is an Education Platform

We provide Free PDF Notes and Videos Classes for Pharmacy Students

Web Site <http://www.fdspharmacy.in/>

You tube <https://www.youtube.com/channel/UC77iEsiuZolU4pB8WAJIR5Q>

What app <https://chat.whatsapp.com/IzSgXtFEvhS4LN5xhUgq5z>

Telegram <https://t.me/+cvxm17xSloA4MjVI>

Face book <https://www.facebook.com/Fdspharmacy-105764311994440/>

E-mail [fdspharmacyinfo@gmail.com](mailto:fdspharmacyinfo@gmail.com)

# Diploma in Pharmacy 1<sup>st</sup> Year

## Pharmacognosy

### Experiment

To perform the morphological study of Ephedra.

#### Aim:

To perform the morphological study of Ephedra.

#### Reference :

Dr. Gupta G.D , Dr. Sharma Shailesh , Kaur Navjit , “Practical Manual of Pharmacognosy” Published by Nirali Prakashan , Pg.No 39 - 42

#### Synonyms

Ma Huang

#### Biological Source :

Ephedra consists of the dried aerial parts of Ephedra gerardiana Wall, Ephedra sinica Stapf, Ephedra equisetina Bunge, Ephedra nebrodensis Tineo and other Ephedra species, belonging to family Ephedraceae.

#### Morphological Characters

This form of evaluation involves a series of tests using the sense organs. The size and morphology of various species differ from each other on following aspects:

- 1) **Ephedra gerardiana:** It has a cylindrical woody stem with a grey or greenish colour. The stems have nodes, internodes, scaly leaves, and terminal buds. The internodes are 3-4 cm apart, and the nodes are covered in scaly leaves. They have a bitter flavour. Within a two-leaved involucre, the plant has stamens and pistils on separate flowers, staminate flowers in catkins with a membranous perianth, and

pistillate flowers terminal on axillary stalks. Fruit is a succulent cone with two carpels each containing a single seed. slender and erect branches, little scale-like leaves articulated and linked at the base forming a sheath.

- 2) **Ephedra sinica:** The stem is 4-7 mm thick, while the branches are 1-2 mm thick. Branch lengths can be up to 30 cm with internodes of 3-6 cm. The colour of the main stem is dark brown. The opposite, decussate, and subulate leaves are 2-4 mm long. The base of the leaf is reddish-brown, the apex is acute and recurved, and the lamina is white. A pair of sheathing leaves found at the nodes, encircles the stem and is attached at the base.
- 3) **Ephedra equisetina:** Stems are 1.5-2 mm thick, woodier, and more branched. Branch lengths range from 25 to 200 cm, with internodes ranging from 1-2.5 cm in length. The exterior surface is smooth and grey to pale green.
- 4) **Ephedra nebrodensis:** The stems are 15 to 35 cm long, 1-2 mm thick, cylindrical, greenish-yellow in colour, with brownish and prominent nodes and a cracked surface that is fibrous in the cortex but includes brownish powdery matter in the pith. The leaves are astringent and bitter in taste, with a brownish to whitish-brown colour, scaly, connate, opposite, and decussate leaves, acute, pleasing, and slightly aromatic odour.




Figure 13: Ephedra Stem

## Result :

The morphological study of Ephedra was performed.

**Hello**

**Friends**

**If you Get Any Help From This Notes /  
Videos**

**Next You Turn To Help Us**

**Please Contribute Some Amount**

**To Our**

**FDSPharmacy Team**

**Phone pe           6398439940**

**Paytm               6398439940**

**Google Pay       6398439940**



Amir Khan

**Thank You**

**Keep Supporting**