

WELCOME

TO



FDSPharmacy
Learn and Educate

This is an Education Platform

We provide Free PDF Notes and Videos Classes for Pharmacy Students

Web Site <http://www.fdspharmacy.in/>

You tube <https://www.youtube.com/channel/UC77iEsiuZolU4pB8WAJIR5Q>

What app <https://chat.whatsapp.com/IzSgXtFEvhS4LN5xhUgq5z>

Telegram <https://t.me/+cvxm17xSloA4MjVI>

Face book <https://www.facebook.com/Fdspharmacy-105764311994440/>

E-mail fdspharmacyinfo@gmail.com

Diploma in Pharmacy 1st Year

Pharmaceutical Chemistry

Experiment

To prepare and submit picric acid from phenol.

Aim:

To prepare and submit picric acid from phenol.

Reference :


‘ Dr. Gupta G.D. , Dr. Sharma Shailish , Kaur Baljeet ’ “Practical Manual of Pharmaceutical Chemistry” Published by Nirali Prakashan, Page no 64 - 68

Requirements:

Erlenmeyer flask (250 mL), volumetric flask, pipette, glass stirring rod, buchner funnel, suction pump, filter papers, measuring cylinder, rubber clock, fume hood, phenol (7.5 ml), nitric acid (22.5 ml), sulphuric acid (20 ml) and cold water (100 ml).

Theory:

Picric acid is a derivative of 2, 4, 6-trinitrophenol. Its molecular formula is $C_6H_3N_3O_7$. It is yellow in colour. Because of the presence of phenol, it is employed as a topical anti-infective and disinfectant, as well as a cleaning agent and preservative. Picric acid is obtained from phenol that has been nitrated. Picric acid is often used to mark rats and mice in pharmacology experiments.





Aromatic hydrocarbons such as 2, 4, 6-trinitrophenol or their substituted derivatives react with conc Nitric acids undergo the nitration reaction (electrophilic substitution reaction) in the presence of strong sulphuric acid (i.e. hydrogen atoms are replaced by nitro group). Even in the absence of sulphuric acid, nitration can be done in some cases.

An important class of organic compound synthesis is the electrophilic aromatic substitution reaction. The position and extent of substitution of new incoming groups are determined by substituents which are already present in the benzene nucleus.

Chemical Reactions

Chemical Reactions



Procedure:

- 1) 7g of phenol and 20 ml of conc. Sulphuric acid should be taken in 500 ml dried round bottomed flask. The mixture should be shaken and heated on a water bath for half an hour.
- 2) A clear solution of o- and p-phenol sulfonic acids should be obtained during this time. The flask should be then cooled in an ice-bath.
- 3) 22.5 ml of conc. HNO₃ should be gradually added with continuous shaking. An exothermic reaction occurs, releasing vast amounts of red fumes (nitrogen oxide) and turning the liquid a deep red colour.
- 4) The flask should be heated in water bath for 2 hour with infrequent shaking. The flask should be cooled and 100 ml of water should be added to it and then cool again.
- 5) The solid should be filtered and washed completely with cold water. The crude picric acid should be recrystallized from aqueous hot alcohol.

Calculation

Here limiting reagent is phenol; hence yield should be calculated from its amount taken.

Molecular formula of phenol = C₆H₆O

Molecular formula of picric acid = C₆H₃N₃O

Molecular weight of phenol = 94 g/mol

Molecular weight of picric acid = 229.10 g/mol

94 g of phenol reacts with benzoyl chloride to give 229.10 gm of picric acid. Therefore, 7.5 of phenol will give x gm of picric acid

$$x = (229.10 \times 7.5) / 94 = 18.27 \text{g}$$

Theoretical yield = 18.27 g

Practical yield = assume 15.3 gm

Percentage (Practical yield x100)

$$= (15.3 \times 100) / 18.27$$

$$83.74\%$$

Uses :

- ✓ It is utilised as a yellow dye for silk and wool, as an explosive as well as an antiseptic in burn therapy.
- ✓ By far, the most common application has been in weaponries and explosives.
- ✓ Explosive D. also called as Dunnite, is an ammonium salt of picric acid that is more powerful than but less stable than TNT (which is produced in a similar process to picric acid but with toluene as the feedstock).
- ✓ Picramide can be further marinated to produce the very stable explosive TATB by aminating picric acid (usually starting with Dunnite)
- ✓ It has been used in organic chemistry to make crystalline salts of organic bases (picrate's) for the sake of identification and characterization.

Result: Picric acid was prepared and submitted. Its practical yield was found to be 15.3 gm and percent yield about 83.74%.

FDSPharmacy
Learn and Educate

Hello

Friends

**If you Get Any Help From This Notes /
Videos**

Next You Turn To Help Us

Please Contribute Some Amount

To Our

FDSPharmacy Team

Phone pe 6398439940

Paytm 6398439940

Google Pay 6398439940



Amir Khan

Thank You

Keep Supporting