WELCOME



This is an Education Platform

We provide Free PDF Notes and Videos Classes for Pharmacy Students

Web Site http://www.fdspharmacy.in/

You tube https://www.youtube.com/channel/UC77iEsiuZolU4pB8WAJIR5Q

What app https://chat.whatsapp.com/IzSgXtFEvhS4LN5xhUgq5z

Telegram https://t.me/+cvxm17xSloA4MjVl

Face book https://www.facebook.com/Fdspharmacy-105764311994440/

E-mail fdspharmacyinfo@gmail.com



Diploma in Pharmacy 2nd Year Community Pharmacy & Management Experiment

To study appropriate handling with correct administration techniques of nasal drops.

Aim:

To study appropriate handling with correct administration techniques of nasal drops.

Reference:

'Dr. Gupta G.D., Dr. Sharma Shailesh, Dr. Gupta Richa, "Practical Manual of Community Pharmacy and Management" Published by Nirali Prakashan, Page no 93 – 96

Theory:


Nasal drops are drugs in solution form which with the help of a dropper or a plastic squeeze bottle are instilled into the nose. Nasal drops are aqueous or oily solutions which are instilled into the nostrils using a dropper. These solutions contain antiseptics, local analgesics, or vasoconstrictors.

Handling and Administration Technique of Nasal Drops

- 1) Nose should be blown gently.
- 2) Hands should be washed completely with soap and water.
- 3) The dropper tip should be examined to confirm if it is damaged or broken
- 4) The dropper tip should be avoided from touching the nose.
- 5) Head should be tilted back during sitting down or lying down on a fly surface with head hanging over the edge
- 6) The medication should be applied in to the nostril as directed
- 7) Head should be bent forward towards right and left consecutively for a few times

Page | 2

- 8) This position should be maintained for a few minutes.
- 9) The cap should be replaced and hands should be washed



Precautions

- 1) This medicine should only be applied into the nostrils 11 should not be taken ocally
- 2) Decongestant-containing nasal drops should not be used continuously for longer than recommendation. Its effectiveness may be lowered by overuse Furthermore, sudden stopping of continuous-long term will result in rebound nasal block.
- 3) This medicine should not be used after 28 days it has been opened.
- 4) Each bottle should be used by one patient to stop the transmission of infection.

Result:

Appropriate handling with correct administration techniques of nasal drops was studied



Hello

Friends

If you Get Any Help From This Notes /

Videos

Next You Turn To Help Us

Please Contribute Some Amount

To Our

FDSPharmacy Team

Phone Pe 6398439940

6398439940 **Paytm**

Google Pay 6398439940



Amir Khan

Thamk You Keep Supporting

