Experiment

WELCOME TO Definition of the second s

This is an Education Platform

We provide Free PDF Notes and Videos Classes for Pharmacy Students

Web Site http://www.fdspharmacy.in/

You tube https://www.youtube.com/channel/UC77iEsiuZolU4pB8WAJIR5Q

What app https://chat.whatsapp.com/IzSgXtFEvhS4LN5xhUgq5z

Telegram <u>https://t.me/+cvxmi7xSloA4MjVl</u>

Face book <u>https://www.facebook.com/Fdspharmacy-105764311994440/</u>

E-mail fdspharmacyinfo@gmail.com



Diploma in Pharmacy 2nd Year Pharmacology Experiment

To study the laboratory animal Rabbit

Aim:

To study the laboratory animal Rabbit

Reference :

[•] Dr. P.Mudagal Manjunatha , "Practical Manual of Pharmacology" Published by Nirali Prakashan, Page no 15 - 18

Theory : Description

- Rectal Temperature: 38.7°C -39.1°C
- Normal Respiratory Rate: 55 per min
- Pulse Rate: 135 per min
- Gestation Period: 28-31 days
- Weaning Age: 6-8 weeks
- Mating Age: 6-9 months
- Litters: 4 Yearly; average 4 litter
- Room Temperature: 15.5-18.5
- Humidity: 40-45%
- Weight (Adult): 0.9-6.75 kg

Housing

- Cages should be made of galvanised iron.
- The minimum size of cage for a medium sized rabbit should be 2x2x1(1/2)ft.
- Young rabbits up to 3 months of age should be housed together, after which males and females should be separated.



Page | 2

Feeding

- → Pelleted diet, such as 18 of Bruce and Parkers(1947) or commercial breeders pellets, should be provided.
- → Daily supply of 72 gm of a mixture of one part oats and three parts bran should be fed as a slightly moist mash.
- \rightarrow Green stuffs or root vegetables should be provided.
- \rightarrow Clean drinking water should be provided.

Handling of Rabbit

- **4** Rabbit's car should be smoothened back
- The cars and loose skin at the back of the neck should be picked up with one hand in a firm grip
- The other hand should be placed under the hind quarter to support the weight and lift gently.
- **4** Rabbits should never be lifted by ear alone.
- **4** Rabbits should be placed on a non-slippery surface.
- If restraint is required during anaesthesia or inoculation, rabbits should be wrapped in a roller towel or placed in a special box.





Common Diseases of Rabbit

- Coccidiosis (hepatic and intestinal)
- Pseudo tuberculosis
- Respiratory infections (Snuffles)
- Pneumococci
- Streptococci
- Intestinal infections (Mucoid enteritis, Dianthoca)
- Rabbit syphilis (Treponema cuniculi)
- Worms (Cysticercus stage of dog tape worm. Taenia pisiformiz

Experimental Procedures on Rabbit

- **4** Antisera
- \rm Anaesthesia
- Scarification
- 4 Subcutaneous inoculation
- **4** Intra-venous inoculation

Applicatio<mark>n</mark>s

- ✓ Pyrogen testing
- ✓ Bioassay of anti-diabetic, curare form drugs and sex hormones
- Screening of agents affecting capillary permeability
- ✓ Drugs used in glaucoma
- ✓ Pharmacokinetic studies
- ✓ Animal of choice for cardiac studies
- ✓ Teratogenicity studies
- ✓ Suitable for researches on reproduction
- Study on miotic and mydriatics

Result :

The laboratory animal rabbit was studied.

- + Intra-peritoneal inoculation
- Intra-cerebral inoculation
- <mark>+</mark> Intra-testicular
- ✤ Ophthalmic
- Collection of blood.



Hello

Friends

If you Get Any Help From This Notes /

Videos

Next You Turn To Help Us

Please Contribute Some Amount

To Our

FDSPharmacy Team

Phone Pe 6398439940

Paytm 6398439940

Google Pay 6398439940



Amir Khan



